What is Semaglutide or Tirzepatide?

This is a class of medication known as Glucagon peptide receptor antagonists or GLP-1 RAs initially approved for Diabetes Mellitus. Several studies including a double blind study in the New England Journal Medicine showed that these medications can reduce body weight by as much as 20%. Because of their positive impact on weight loss, GLP-1 RAs have been approved by the FDA for weight loss and obesity management.

How is Semaglutide or Tirzepatide supplied?

Tirzepatide or Semaglutide is supplied in a kit that contains a vial of medication with syringes and alcohol pads.

How does Semaglutide work in weight loss

- Semaglutide helps you lose weight by mimicking the effects of the hormone GLP-1 which plays a role in regulating food intake and reducing appetite.
- Semaglutide regulates food intake by increasing the stomach emptying time.
- Semaglutide increases insulin production which results in improved blood sugar levels.
- Semaglutide sends signals to the brain letting your brain know that your stomach is full.

How do I get Semaglutide or Tirzepatide?

Semaglutide and Tirzepatide are prescription medications therefore you need to ask your doctor if you are a good candidate for these medications.

Alternatively:

- Get in touch with us:
 - 🖂 qj@stroudcompounding.com
 - 570-476-6936
 - (sms) 570-828-3643

Directions

 Wash your hands and take the vial from the freezer.
Allow the vial a few minutes to thaw out before you wipe the rubber enclosure with an alcohol pad provided. Allow the vial to air dry.

3. Remove the syringes from their protective covering.

- 4. Remove the needle cap and pull the plunger to the desired dosage to fill the syringe with air.
- 5. Insert the needle into the vial and push the air that was drawn back into the vial

6. Turn the vial upside down to see the needle lines clearly and slowly pull the syringe plunger until the solution reaches the correct dose

Administering the solution

1. This medicine can be injected as a shot under the skin of your stomach, thighs, or upper arm.

2. Use the swab provided to clean the area you want to inject the medication.

3. Use your dominant hand to hold the syringe like a dart with the barrel between your fingers and your thumb. Pinch an area of your skin with your other hand to create a lump.

4. Insert the needle with a quick jab into the pinched skin.

5. Press the plunger down with your pointer finger until the syringe is empty.

6. Pull out the needle at the same angle it went in.

7. Dispose of the used syringe in a sharps container. Do not reuse any syringes.

Warnings and Precautions:

1. Follow the dosage regimen as prescribed by your doctor.

2. Rotate injection sites every week between

your stomach, thighs or upper arms.

3. Do not use more than once a week and

discard vial after 28 days.

4. If you experience any side-effects, please call

your doctor as soon as possible.

Storage Directions:

 $\boldsymbol{\cdot}$ Medication must be stored in an amber vial or

brown bag in the freezer

After injecting yourself, return the vial to the

brown bag or vial provided and put it in the freezer.

· If you leave your medication vial outside the

freezer, discard the vial and syringe

Let's Chat!

gj@stroudcompounding.com

570-476-6936

570-828-3643



1619 N 9th Street Suite 3, Stroudsburg, PA 18360

Phone:(570) 476-6936 Fax: (570) 476-6938

www.stroudcompounding.com